Linking Environmental Risk Factors to Maternal Health Outcomes and Neonatal Wellbeing

Dr. R. Niveditha
Assistant Professor, Department of Obstetrics and Gynecology Nursing, NRI College of Nursing, Guntur, Andhra Pradesh, India.

Received: 12-06-2024; Revised: 22-07-2024; Accepted: 05-08-2024; Published: 22-08-2024

Abstract
Environmental exposure to pollutants and hazardous substances has significant implications for maternal morbidity and mortality, as well as neonatal health. Pregnant women exposed to high levels of air pollution, toxic chemicals, and other environmental hazards are at increased risk of complications such as preeclampsia, preterm birth, and low birth weight. These exposures can lead to long-term health issues for both mothers and their infants, including respiratory problems, developmental delays, and increased susceptibility to chronic diseases. Understanding the pathways through which environmental factors impact maternal and neonatal outcomes is crucial for developing effective public health policies and interventions to mitigate these risks.

Keywords: ***Environmental exposure, Maternal morbidity, Maternal mortality, Neonatal health, Air pollution, Toxic chemicals, Preeclampsia, Preterm birth.***

1.Introduction

Environmental exposure to pollutants and hazardous substances represents a critical and growing concern in public health, particularly regarding its impact on maternal and neonatal health outcomes. As industrial activities, urbanization, and the use of various chemicals proliferate, the prevalence of environmental pollutants such as air pollution, heavy metals, pesticides, and endocrine-disrupting chemicals (EDCs) has escalated. These pollutants can have profound adverse effects on pregnant women and their developing fetuses, potentially leading to severe health complications and increased healthcare burdens(1).

Air pollution, one of the most pervasive environmental hazards, primarily consists of particulate matter (PM), nitrogen dioxide (NO2), sulfur dioxide (SO2), carbon monoxide (CO), and ozone (O3). These pollutants originate from vehicular emissions, industrial discharges, agricultural practices, and natural sources like wildfires. Heavy metals such as lead, mercury, and arsenic are commonly found in contaminated water, soil, and food, often resulting from mining, industrial processes, and improper waste disposal. Pesticides and EDCs, prevalent in agricultural and household products, can persist in the environment and bioaccumulate, posing significant risks to human health.

Exposure to these environmental pollutants during pregnancy has been linked to a range of maternal health issues. Pregnant women exposed to high levels of air pollution, for instance, have an increased risk of developing conditions such as preeclampsia, gestational hypertension, and gestational diabetes. These conditions not only complicate pregnancy but also contribute to maternal morbidity and mortality. The underlying mechanisms are thought to involve systemic inflammation, oxidative stress, and disruptions in placental function, which collectively impair maternal health(2).

The health implications for neonates are equally concerning. Fetuses exposed to environmental pollutants in utero are at greater risk for preterm birth, low birth weight, and intrauterine growth restriction (IUGR). These adverse birth outcomes are significant predictors of neonatal morbidity and mortality, as well as long-term health challenges including respiratory disorders, neurodevelopmental delays, and increased susceptibility to chronic diseases such as asthma and cardiovascular conditions(3). The fetal origins of adult disease hypothesis suggests that environmental exposures during critical periods of development can have lasting impacts on health trajectories.

The biological mechanisms through which environmental pollutants exert their effects involve complex interactions at the molecular, cellular, and systemic levels. For instance, fine particulate matter (PM2.5) can penetrate deep into the respiratory tract and enter the bloodstream, leading to systemic inflammation and oxidative stress. These processes can disrupt placental function, leading to impaired nutrient and oxygen transport to the fetus. Additionally, exposure

to heavy metals and EDCs can interfere with hormonal regulation and epigenetic modifications, further affecting fetal development and maternal health(4).
Addressing the impact of environmental exposure on maternal and neonatal health necessitates comprehensive public health strategies. These strategies should include stringent regulations to limit emissions of harmful pollutants, widespread monitoring of environmental quality, and targeted interventions for high-risk populations. Public health campaigns to raise awareness about the sources and risks of environmental pollutants, along with policies to reduce exposure, are crucial. Furthermore, healthcare providers should be equipped with the knowledge and tools to identify and manage environmental risk factors in pregnant women.

2.Literature Survey

Numerous studies have investigated the relationship between environmental exposure and adverse maternal health outcomes. Research has consistently shown that exposure to air pollution, particularly fine particulate matter (PM2.5), is associated with an increased risk of hypertensive disorders in pregnancy, such as preeclampsia and gestational hypertension(5). A systematic review by Pedersen et al. (2014) found that exposure to PM2.5 during pregnancy was significantly correlated with elevated blood pressure and an increased likelihood of preeclampsia. Similarly, a cohort study by Dadvand et al. (2013) indicated that pregnant women living in areas with high levels of NO2 had a higher incidence of gestational diabetes, suggesting a link between air quality and metabolic health during pregnancy(6).
The impact of heavy metals on maternal health has also been well-documented. Lead exposure, for example, has been shown to be detrimental to both maternal and fetal health. A meta-analysis by Liu et al. (2018) demonstrated that elevated blood lead levels in pregnant women were associated with a higher risk of spontaneous abortion, preterm birth, and low birth weight. Arsenic exposure, often through contaminated drinking water, has been linked to increased rates of stillbirth and neonatal mortality, as reviewed by Quansah et al. (2015). These findings underscore the critical need for monitoring and mitigating heavy metal exposure in vulnerable populations(7).
Neonatal health outcomes are profoundly affected by environmental exposures as well. The literature indicates that prenatal exposure to air pollution can lead to adverse birth outcomes such as preterm birth and low birth weight. A comprehensive review by Stieb et al. (2012) highlighted that increased levels of ambient air pollutants were significantly associated with a reduction in birth weight and an increase in preterm deliveries. Additionally, a study by Vrijheid et al. (2011) on the effects of exposure to industrial pollutants found similar outcomes, with higher incidences of low birth weight and congenital anomalies in neonates.
Endocrine-disrupting chemicals (EDCs) are another critical area of concern. These chemicals, found in many household and industrial products, can interfere with hormone function and have been linked to developmental and reproductive health issues. A study by Ferguson et al. (2014) reported that higher urinary concentrations of phthalates, a common EDC, during pregnancy were associated with an increased risk of preterm birth. Another study by Braun et al. (2016) found that prenatal exposure to bisphenol A (BPA) was linked to neurobehavioral issues in children, suggesting long-term developmental impacts.
The mechanisms underlying these associations are complex and multifaceted. Oxidative stress and inflammation are common pathways through which pollutants exert their effects. For instance, Shah et al. (2011) discussed how air pollution-induced oxidative stress could impair placental function, leading to adverse birth outcomes. Epigenetic modifications, as reviewed by Breton et al. (2017), also play a crucial role, with environmental exposures potentially altering gene expression and impacting fetal development.
In summary, the literature clearly indicates that environmental exposures have significant and multifaceted impacts on maternal morbidity and mortality, as well as neonatal health. These findings highlight the urgent need for targeted public health interventions, improved regulatory policies, and continued research to better understand and mitigate these risks(8).

3.Existing and Proposed System

Current systems addressing the impact of environmental exposure on maternal and neonatal health include regulatory frameworks, monitoring programs, and public health initiatives. Regulatory bodies such as the Environmental Protection Agency (EPA) and the World Health Organization (WHO) have established guidelines and standards for air and water quality to limit exposure to harmful pollutants. For instance, the Clean Air Act in the United States mandates limits on emissions of hazardous air pollutants from industrial sources. Monitoring programs, like the Air Quality Index (AQI) and various biomonitoring initiatives, provide critical data on environmental contaminant levels and exposure trends. Public health initiatives often focus on education and outreach, informing pregnant women and healthcare providers about the risks of environmental exposures and ways to minimize them.

Despite these efforts, significant gaps remain in protecting maternal and neonatal health from environmental hazards. Proposed systems aim to address these gaps through enhanced regulatory measures, comprehensive environmental health surveillance, and targeted interventions. Strengthening regulations to reduce permissible levels of pollutants, particularly in areas with vulnerable populations, is crucial. Expanding monitoring efforts to include a wider array of contaminants and more frequent sampling can provide better data for risk assessment and policy-making. Additionally, integrating environmental health considerations into prenatal care programs can improve early detection and management of exposure risks. Proposed systems also emphasize the importance of multidisciplinary research to uncover the mechanisms of pollutant effects and the development of innovative technologies for pollution control and exposure reduction. By adopting these comprehensive and proactive approaches, we can better safeguard maternal and neonatal health against the adverse effects of environmental exposure.

4.Current State of Maternal Morbidity, Mortality, and Neonatal Health Worldwide

The health of mothers and newborns is a crucial indicator of societal well-being and development. Despite significant progress in healthcare over the past few decades, maternal morbidity and mortality, along with neonatal health challenges, remain significant global public health issues, particularly in low- and middle-income countries (LMICs).

Maternal Morbidity and Mortality

Maternal morbidity refers to health complications experienced by women during pregnancy, childbirth, or the postpartum period. These complications can have long-term health impacts and often lead to maternal mortality if not properly managed. The World Health Organization (WHO) estimates that approximately 295,000 women died during and following pregnancy and childbirth in 2017, with the vast majority of these deaths occurring in LMICs. Sub-Saharan Africa and South Asia bear the highest burdens, accounting for about 86% of all maternal deaths globally.

Several factors contribute to high maternal morbidity and mortality rates, including limited access to quality healthcare, inadequate infrastructure, and socio-economic barriers. Common causes of maternal mortality include severe bleeding (often postpartum hemorrhage), infections, high blood pressure during pregnancy (pre-eclampsia and eclampsia), complications from delivery, and unsafe abortions. Addressing these issues requires comprehensive healthcare strategies that encompass preventive, diagnostic, and therapeutic measures(9).

Neonatal Health

Neonatal health, referring to the health of infants within the first 28 days of life, is a critical component of child health. According to UNICEF, neonatal deaths accounted for 47% of all under-five child deaths in 2019, with approximately 2.4 million neonatal deaths occurring that year. The leading causes of neonatal deaths include preterm birth complications, intrapartum-related events (such as birth asphyxia), infections (including sepsis, pneumonia, and tetanus), and congenital anomalies.

Neonatal mortality rates are highest in regions with limited healthcare access and resources. For instance, in sub-Saharan Africa, the neonatal mortality rate was 27 deaths per 1,000 live births in 2019, compared to 3 deaths per 1,000 live births in high-income countries. Effective neonatal health interventions include providing skilled birth attendance, improving prenatal and postnatal care, promoting breastfeeding, and ensuring access to essential medicines and vaccines.

Global Initiatives and Progress

Global health initiatives have made considerable strides in addressing maternal and neonatal health challenges. The United Nations' Sustainable Development Goals (SDGs), particularly SDG 3, aim to reduce the global maternal mortality ratio to less than 70 per 100,000 live births and end preventable deaths of newborns and children under five by 2030. Programs such as the WHO's Every Woman Every Child initiative and the Partnership for Maternal, Newborn & Child Health (PMNCH) have been pivotal in mobilizing resources and implementing strategies to improve health outcomes(10).

Efforts to enhance maternal and neonatal health have also focused on improving healthcare infrastructure, increasing the availability of skilled health professionals, and expanding access to essential services. For example, increasing the number of healthcare facilities offering comprehensive emergency obstetric and neonatal care (CEmONC) has been a key strategy in reducing maternal and neonatal mortality. Additionally, community-based interventions, such as training traditional birth attendants and community health workers, have proven effective in regions where access to formal healthcare is limited.

Challenges and Future Directions

Despite progress, significant challenges remain in achieving global maternal and neonatal health goals. Socio-economic inequalities, political instability, and cultural barriers continue to hinder access to quality healthcare. Additionally, emerging health threats such as infectious diseases (e.g., COVID-19) and climate change pose new risks to maternal and neonatal health(11).

Future efforts must prioritize the integration of maternal and neonatal health services into broader health systems, ensuring that these services are sustainable and resilient. Innovations in healthcare delivery, such as telemedicine and mobile health (mHealth) technologies, can play a crucial role in reaching underserved populations. Moreover, increasing investments in health systems strengthening, education, and community engagement is essential to address the root causes of maternal and neonatal health disparities.

5.Cost of Maternal Illness and Death in the States of India.

India faces a significant burden of maternal morbidity and mortality, with stark disparities across its states. Although the country has made notable progress in reducing maternal mortality over the past few decades, it continues to grapple with high rates of maternal health complications and deaths, particularly in poorer and rural regions. According to the Sample Registration System (SRS) data from 2016-18, India's Maternal Mortality Ratio (MMR) was 113 per 100,000 live births, a substantial decline from previous years but still above the global target set by the Sustainable Development Goals (SDGs).

Regional Disparities

The distribution of maternal mortality is uneven across Indian states, reflecting varying levels of healthcare access, socio-economic development, and infrastructural support. States like Kerala, Tamil Nadu, and Maharashtra have achieved relatively low MMRs, often cited as examples of effective health systems and policies. For instance, Kerala reported an MMR of 43 per 100,000 live births, owing to its robust healthcare infrastructure, high literacy rates, and effective maternal health programs. In contrast, states such as Uttar Pradesh, Bihar, and Madhya Pradesh struggle with high MMRs, with figures exceeding 150 per 100,000 live births in some areas(12).

Contributing Factors

Several factors contribute to the high maternal morbidity and mortality in these states. Limited access to quality healthcare services is a primary concern. In rural and remote areas, women often lack access to skilled birth attendants and emergency obstetric care. This gap is exacerbated by a shortage of healthcare facilities and trained personnel. Additionally, socio-economic barriers, including poverty, low literacy levels, and gender inequality, significantly impede women's access to necessary health services. Cultural practices and a lack of awareness about maternal health further compound these challenges, leading to delays in seeking and receiving appropriate care.

Healthcare Infrastructure and Services

The disparity in healthcare infrastructure between states plays a crucial role in maternal health outcomes. States with lower MMRs have better healthcare infrastructure, including well-equipped hospitals, availability of skilled healthcare providers, and effective referral systems. In contrast, high-MMR states often suffer from inadequate healthcare facilities, poorly maintained medical equipment, and insufficient numbers of trained health workers. These issues are particularly acute in rural and tribal areas, where healthcare access is severely limited. Efforts to improve healthcare infrastructure and services in high-MMR states are critical to addressing maternal health disparities(13).

Government Initiatives and Interventions

The Indian government has launched several initiatives aimed at reducing maternal morbidity and mortality. Programs such as the Janani Suraksha Yojana (JSY) and the Janani Shishu Suraksha Karyakram (JSSK) have been instrumental in promoting institutional deliveries and providing free maternal and child health services. The JSY scheme, for instance, offers financial incentives to encourage institutional deliveries, particularly among low-income families. The JSSK program ensures free transportation, medicines, and nutritional support for pregnant women and newborns. These initiatives have contributed to increasing institutional delivery rates and improving maternal health outcomes in many states.

However, the implementation and impact of these programs vary across regions. States with effective administrative and health systems have seen more substantial benefits from these initiatives, while others lag due to poor implementation and monitoring. Strengthening the delivery and oversight of maternal health programs is essential to ensure that the benefits reach the most vulnerable populations(14).

Community-Based Interventions

Community-based interventions have also shown promise in reducing maternal morbidity and mortality in high-burden states. Training and deploying Accredited Social Health Activists (ASHAs) and other community health workers have been effective in promoting maternal health awareness and facilitating access to healthcare services. These workers play a crucial role in educating women about antenatal care, institutional deliveries, and postpartum care, as well as assisting with referrals and transportation to health facilities.

6. Different environmental factors that affect maternal health

Environmental factors play a crucial role in maternal health, but there is a significant disparity in exposure and impact across different regions and socio-economic groups. Environmental pollutants such as air and water pollution, toxic chemicals, and inadequate sanitation facilities disproportionately affect vulnerable populations, exacerbating maternal morbidity and mortality. The burden of these environmental risks is particularly pronounced in low- and middle-income countries (LMICs) and within marginalized communities, highlighting the intersection of environmental justice and public health.

Air Pollution and Maternal Health

Air pollution is a pervasive environmental hazard that has profound effects on maternal health. Exposure to high levels of particulate matter (PM2.5 and PM10), nitrogen dioxide (NO2), sulfur dioxide (SO2), and other pollutants has been linked to adverse pregnancy outcomes such as preterm birth, low birth weight, and preeclampsia. Studies have shown that pregnant women in urban areas with heavy traffic and industrial activities are at higher risk due to increased exposure to these pollutants. For instance, research conducted in India revealed that pregnant women exposed to high levels of air pollution in cities like Delhi are more likely to experience adverse pregnancy outcomes compared to those in rural or less polluted areas. This disparity is often exacerbated by socio-economic factors, as poorer populations tend to live in areas with higher pollution levels due to cheaper housing and proximity to industrial zones.

Water Pollution and Sanitation

Access to clean water and proper sanitation is another critical environmental determinant of maternal health. Contaminated water sources, often polluted with pathogens, heavy metals, and industrial chemicals, pose significant health risks. Inadequate sanitation facilities increase the likelihood of waterborne diseases such as cholera and diarrhea, which can be particularly dangerous during pregnancy. Rural and impoverished regions are more affected by these issues due to the lack of infrastructure and resources. For example, in many parts of sub-Saharan Africa and South Asia, women are more likely to suffer from water-related health issues due to reliance on unsafe water sources and inadequate sanitation. These conditions contribute to maternal infections, complications during childbirth, and higher mortality rates.

Chemical Exposures

Exposure to toxic chemicals, including pesticides, heavy metals, and endocrine-disrupting chemicals (EDCs), is another environmental risk factor with significant implications for maternal health. Agricultural workers and those living near industrial sites are particularly vulnerable. For instance, in agricultural regions of Latin America and Asia, women are often exposed to pesticides, which have been associated with miscarriages, stillbirths, and congenital anomalies. Similarly, industrial activities can lead to elevated levels of heavy metals like lead and mercury in the environment, which are known to cause severe reproductive and developmental health issues. These exposures are often higher in economically disadvantaged areas where regulatory oversight may be weaker, and residents may lack the resources to mitigate exposure risks.

Indoor Air Pollution

Indoor air pollution, primarily from the use of biomass fuels for cooking and heating, is a significant environmental health issue in many developing countries. Women, who spend a considerable amount of time near cooking stoves, are at a higher risk of inhaling harmful pollutants such as carbon monoxide and fine particulate matter. This exposure is linked to respiratory infections, low birth weight, and other adverse pregnancy outcomes. The disparity in indoor air pollution exposure is stark, with rural households in LMICs being the most affected due to the continued reliance on traditional cooking methods and lack of access to clean energy alternatives.

Socio-Economic Disparities and Policy Gaps

The disparity in environmental causes for maternal health is closely tied to socio-economic inequalities and policy gaps. Wealthier populations have better access to clean environments, healthcare, and resources to mitigate environmental risks. In contrast, poorer and marginalized communities are often left to contend with higher levels of pollution and fewer resources to address health issues. Policy gaps at both national and local levels further exacerbate these disparities. In many LMICs, environmental regulations may be inadequately enforced, and health policies may not sufficiently address the unique needs of pregnant women in polluted environments. Additionally, there is often a lack of targeted public health interventions and resources to support the most vulnerable populations.

7. Health issues and typical pregnancy-related risk factors

Pregnancy, while a natural and often joyous process, can present a range of medical conditions and risk factors that affect both the mother and the fetus. Understanding these conditions and risk factors is crucial for ensuring the health and safety of pregnant women and their babies. Medical conditions during pregnancy can vary widely in severity and impact, from mild and manageable to severe and life-threatening.

Hypertensive Disorders of Pregnancy

One of the most significant medical conditions associated with pregnancy is hypertensive disorders, which include chronic hypertension, gestational hypertension, preeclampsia, and eclampsia. Chronic hypertension refers to high blood pressure present before pregnancy or diagnosed before the 20th week of gestation. Gestational hypertension, on the other hand, develops after the 20th week and can progress to preeclampsia if left unmanaged. Preeclampsia is characterized by high blood pressure and signs of damage to another organ system, often the kidneys, evidenced by proteinuria. It can lead to serious complications, including fetal growth restriction, placental abruption, and organ failure in the mother. Eclampsia is a severe form of preeclampsia that includes seizures and can be fatal for both mother and child if not promptly treated.

Gestational Diabetes Mellitus (GDM)

Gestational diabetes mellitus (GDM) is another common medical condition that can arise during pregnancy. It involves high blood sugar levels that develop during pregnancy and usually resolve after delivery. GDM can pose significant risks to both the mother and the fetus, including increased risk of cesarean delivery, macrosomia (large birth weight), and neonatal hypoglycemia. Women with GDM are also at a higher risk of developing type 2 diabetes later in life. Risk factors for GDM include obesity, a family history of diabetes, previous history of GDM, and certain ethnic backgrounds such as South Asian, African American, Hispanic, and Native American.

Infections During Pregnancy

Infections during pregnancy can lead to complications and pose risks to both the mother and the developing fetus. Common infections include urinary tract infections (UTIs), bacterial vaginosis, and sexually transmitted infections (STIs) such as chlamydia, gonorrhea, and syphilis. More severe infections like toxoplasmosis, rubella, cytomegalovirus (CMV), and Zika virus can cause congenital infections, leading to serious birth defects and developmental issues. Pregnant women are also at higher risk for complications from influenza and COVID-19, highlighting the importance of vaccination and preventive measures.

Preterm Birth and Premature Rupture of Membranes (PROM)

Preterm birth, defined as delivery before 37 weeks of gestation, is a leading cause of neonatal morbidity and mortality. Various risk factors contribute to preterm birth, including a history of preterm delivery, multiple pregnancies (twins, triplets), infections, and conditions such as cervical insufficiency. Premature rupture of membranes (PROM), where the amniotic sac ruptures before labor begins, can also lead to preterm birth and increases the risk of infection for both the mother and the baby. Management of preterm birth involves monitoring and interventions to delay delivery, if possible, and ensuring the availability of neonatal intensive care for premature infants.

Placental Complications

Placental complications such as placenta previa and placental abruption are serious conditions that can threaten maternal and fetal health. Placenta previa occurs when the placenta covers the cervix, causing severe bleeding during pregnancy and delivery. Placental abruption, where the placenta detaches from the uterine wall before delivery, can lead to heavy bleeding, fetal distress, and preterm birth. Both conditions require careful monitoring and often necessitate early delivery via cesarean section to prevent severe complications.

Maternal Age and Pregnancy

Maternal age is a significant risk factor for various pregnancy complications. Advanced maternal age (35 years and older) is associated with increased risks of chromosomal abnormalities such as Down syndrome, miscarriage, stillbirth, and complications like preeclampsia and GDM. Younger maternal age (teenage pregnancies) is also linked to higher rates of preterm birth, low birth weight, and neonatal complications. These age-related risks underscore the importance of preconception counseling and appropriate prenatal care to manage potential complications.

Lifestyle Factors and Pregnancy

Lifestyle factors, including diet, physical activity, and substance use, significantly impact pregnancy outcomes. Poor nutrition and obesity are associated with gestational diabetes, hypertensive disorders, and complications during labor and delivery. Conversely, undernutrition can lead to fetal growth restriction and low birth weight. Physical activity, while generally beneficial, should be moderated according to medical advice during pregnancy. Substance use, including smoking, alcohol consumption, and illicit drug use, poses severe risks to fetal development and can lead to miscarriage, stillbirth, preterm birth, and congenital abnormalities. Pregnant women should receive counseling on healthy lifestyle choices to minimize these risks.

Mental Health Conditions

Mental health conditions, such as depression and anxiety, are prevalent during pregnancy and the postpartum period. These conditions can affect maternal well-being and pregnancy outcomes, including preterm birth and low birth weight. Untreated mental health issues can also impair maternal-infant bonding and increase the risk of postpartum depression. It is crucial to screen for and address mental health concerns as part of comprehensive prenatal care.

Managing medical conditions and mitigating risk factors associated with pregnancy is essential for the health of both the mother and the fetus. Comprehensive prenatal care, which includes regular monitoring, appropriate interventions, and patient education, plays a critical role in identifying and managing these conditions. By addressing the diverse factors that influence pregnancy outcomes, healthcare providers can help ensure safer pregnancies and healthier babies.

8.Conclusion and Future work

Environmental exposures significantly impact maternal morbidity and mortality and neonatal health, posing serious public health challenges, particularly in low- and middle-income countries (LMICs) and marginalized communities. Exposure to air and water pollution, toxic chemicals, inadequate sanitation, and indoor pollutants like biomass fuels can lead to severe pregnancy complications such as preeclampsia, gestational diabetes, preterm birth, low birth weight, and congenital anomalies. These environmental risks are compounded by socio-economic disparities, limited access to quality healthcare, and insufficient regulatory measures.

To mitigate the adverse effects of environmental exposures on maternal and neonatal health, a multi-faceted approach is required. This includes strengthening regulatory frameworks to reduce pollution levels, improving healthcare infrastructure, and ensuring equitable access to clean environments and healthcare services. Targeted public health interventions, such as promoting the use of clean cooking technologies and enhancing water and sanitation facilities, are essential. Additionally, increasing awareness and education about environmental risks and their impact on health can empower communities to take preventive measures.

Future Work

1. Enhanced Monitoring and Research:

Future work should focus on enhancing environmental health surveillance and research. This includes comprehensive monitoring of air and water quality, as well as the identification and quantification of toxic chemicals in various environments. Longitudinal studies examining the effects of specific pollutants on maternal and neonatal health outcomes are crucial. This research can provide valuable data to inform policy-making and the development of targeted interventions.

2. Strengthening Healthcare Systems:

Improving healthcare infrastructure and services in regions with high maternal morbidity and mortality is critical. This includes increasing the availability of skilled healthcare providers, ensuring access to emergency obstetric care, and integrating environmental health considerations into maternal and neonatal healthcare programs. Strengthening healthcare systems also involves training healthcare workers to recognize and manage health issues related to environmental exposures.

3. Policy and Regulation:
Strengthening and enforcing environmental regulations is vital to reducing pollution levels and protecting vulnerable populations. Governments should implement stricter standards for industrial emissions, vehicle emissions, and waste management. Policies should also promote the use of clean energy sources and sustainable agricultural practices to minimize environmental contamination. Ensuring that these regulations are effectively enforced is crucial for their success.
4. Community Engagement and Education:
Community-based interventions and education programs can play a significant role in mitigating environmental risks. Engaging communities in discussions about environmental health risks and preventive measures can enhance local capacity to address these issues. Educational campaigns should focus on promoting safe practices, such as the use of clean cooking technologies, proper sanitation, and safe water storage.
5. Technology and Innovation:
Investing in technological innovations can help reduce environmental exposures and improve health outcomes. This includes developing affordable clean energy technologies, improving water purification systems, and creating low-cost pollution monitoring devices. Innovative solutions that are accessible and scalable in LMICs can have a significant impact on reducing environmental health risks.
6. Global Collaboration:
Addressing the global burden of environmental exposures on maternal and neonatal health requires international collaboration. Sharing knowledge, resources, and best practices can help countries implement effective strategies and interventions. Global health organizations, governments, and non-governmental organizations (NGOs) should work together to address these challenges and support research, policy development, and program implementation.
In conclusion, reducing the impact of environmental exposures on maternal morbidity and mortality and neonatal health requires a coordinated effort that spans research, policy, healthcare, and community engagement. By prioritizing these areas in future work, we can move towards a world where all women and children have the opportunity to live in safe and healthy environments

Acknowledgement: Nil

Conflicts of interest
The authors have no conflicts of interest to declare

References

1. Sharma RS, Mathur VK. Impact of Air Pollution on Pregnancy Outcomes in Urban India. Indian Journal of Public Health. 2016; 60(4): 287-294.
2. Patel D, Desai P. Heavy Metal Exposure and Pregnancy Complications: A Study from Gujarat, India. Journal of Environmental Biology. 2017; 38(5): 913-918.
3. Singh P, Agarwal A. Maternal Exposure to Pesticides and Adverse Birth Outcomes in Rural India. Environmental Science and Pollution Research. 2018; 25(23): 22972-22979.
4. Kumar A, Gupta A. Climate Change and Maternal Health: Evidence from India. Indian Journal of Medical Research. 2019; 149(6): 731-737.
5. Reddy KS, Prasad JV. Noise Pollution and Pregnancy: An Indian Perspective. Indian Journal of Community Medicine. 2020; 45(3): 279-285.
6. Sinha R, Jha S. Impact of Endocrine Disruptors on Pregnancy Outcomes in Indian Women. Journal of Reproductive Health and Medicine. 2021; 7(4): 191-198.
7. Choudhary R, Singh M. Socioeconomic Disparities and Environmental Exposures During Pregnancy in India. Indian Journal of Occupational and Environmental Medicine. 2018; 22(3): 138-143.

8. Rao V, Prakash S. Airborne Pollutants and Neonatal Health: Insights from Indian Studies. Journal of Environmental Health Science and Engineering. 2016; 14: 22.
9. Nair S, Raju R. Prenatal Environmental Stressors and Neurodevelopment in Indian Infants. Indian Journal of Pediatrics. 2017; 84(9): 697-704.
10. Kumar V, Bansal A. Hazardous Chemical Exposure and Pregnancy Outcomes: Evidence from India. Journal of Occupational Health. 2018; 60(5): 382-389.
11. Sharma M, Singh S. Environmental Justice and Birth Outcomes in India. Indian Journal of Social Work. 2019; 80(2): 175-182.
12. Gupta R, Ghosh S. Urban Pollution and Maternal Health in India: A Review. Journal of Environmental Management. 2020; 253: 109-748.
13. Tripathi S, Mishra R. Proximity to Industrial Sites and Pregnancy Risks in Indian Women. Science of The Total Environment. 2021; 763: 142-952.
14. Raj A, Shankar V. Green Space and Pregnancy Outcomes: A Study from Urban India. Environmental Pollution. 2022; 292: 118-435.